

RBE, determined at higher dose levels for postradiation alopecia. Mitotic indices were depressed for a 3-fold longer interval than that observed after a "comparable" dose of x-rays. Marked alterations in labeling indices, probably reflecting prolonged cell retention and retarded maturation rates, and discrete reductions in matrix cell populations characterized postneutron radiation changes and were entirely absent in the x-ray study. At the cellular level, these findings support the results of previous investigations [1,9] which have recorded considerably higher RBE's for neutron radiation in very low dose ranges compared to those found at higher dose levels. It should be emphasized that our current study comprises a single low-dose experiment. Although a more complete dose response curve to determine corresponding variations in RBE would be a laborious project, the resulting data would be of considerable interest.

Access to the fast neutron facility at the Franklin McLean Memorial Research Institute, the Pritzker School of Medicine, was kindly provided by Drs. M.L. Griem and F.T. Kuchnir.

REFERENCES

1. Field SB: An historical survey of radiobiology and radiotherapy with fast neutrons. *Curr Topics Rad Res Qtrly* 11:1-86, 1976
2. Hornsey S., Andreozzi U, Warren PR: Sublethal damage in cells of the mouse gut after mixed treatment with x-rays and fast neutrons. *Br J Radiol* 50:513-517, 1977
3. Catterall M, Southerland I, Bewley DK: First results of a randomised clinical trial of fast neutrons compared with X or gamma rays in the treatment of advanced tumours of the head and neck. *Br Med J* 2:653-656, 1975
4. Parker RG, Berry HC, Caderao JB, Gerdes AJ, Hussey DH, Ornitz R, Rogers CC: Preliminary clinical results from U.S. fast neutron teletherapy studies. *Cancer* 40:1434-1438, 1977
5. Malkinson FD, Griem ML, Marianovic R: Follicle squash preparations: uses in studies of cell kinetics following irradiation. *J Invest Dermatol* 57:382-388, 1971
6. Skaggs LS, Kuchnir FT, Harper PV, Griem ML: Physical characteristics of fast-neutron beams from a compact medical cyclotron. *Radiol* 111:471-472, 1974
7. Fry RJM, Kessler D, Kisieleski W, Weber C, Griem ML, Malkinson FD: A method for the study of cell proliferation of the hair follicle. Argonne Natl Laboratory Biological and Medical Research Division Annual Report, ANL-7535, 1968
8. Malkinson FD, Keane JT: Hair matrix cell kinetics: a selective review. *Int J Dermatol* 17:536-551, 1978
9. Chow DC, Rothman S, Lorincz AL, Malkinson F, Sandburg A: Reduction in rate of growth of hair in mice as an indicator of exposure to chronic low dosage ionizing radiation. *Nature* 203: 847-848, 1964

Gordon Research Conference on Epithelial Differentiation and Keratinization

The first Gordon Research Conference directed specifically at problems of epithelial differentiation and keratinization will be held on August 6 through 10, 1979 at the Tilton School in Tilton, New Hampshire. This conference will be chaired by Drs. Irwin M. Freedberg and Isadore A. Bernstein and will deal with epithelial structure, models of epithelial differentiation, normal and pathological epithelial differentiation in vivo and in vitro, physical and chemical epithelial carcinogenesis, repair, epithelial and mesenchymal interactions, and biosynthesis of epithelial proteins and keratin.

Participation in the conference is limited and participants will be selected from applications for attendance. Applications may be obtained from Dr. Alexander M. Cruikshank, Director, Gordon Research Conference, Pastore Chemical Laboratory, University of Rhode Island, Kingston, Rhode Island, 02881.